REG. NO: D7635654-AFINJ JOHP-ISSN: 2348-7704



## ACUTE ORAL TOXICITY STUDY OF METHANOLIC EXTRACT OF ALPINIA OFFICINARUM IN ALBINO WISTER RATS

JOHP

## Deorao Madhaorao Awari <sup>1</sup>, Rahul Somani<sup>2</sup>

- 1. JSPM's Charak College of Pharmacy and Research ,Wagholi, Pune. Savitribai Phule Pune University, Pune.
- 2. Smt.Kashibai Navale College of Pharmacy, Kondhwa, Pune, Savitribai Phule Pune University,Pune

Address for Correspondence: editorjohp@gmail.com

## **ABSTRACT:**

**Objective**: The present study was designed to find out the effective dose of methanolic extract of *Alpinia officinarum Hanse* in female Albino Wistar rats by using OECD guideline 425.

Materials and Method: Adult female albino Wistar rat (150-200 g) were subjected to acute oral toxicity studies as per guideline (AOT no. 425) suggested by Organization for Economic Cooperation and Development (OECD) (2001). Dose of 175, 550, 1750 and 2000 mg/kg of extracts were administered orally to female Wistar rat. Rat was then observed for incidence of mortality or any sign of toxicity for 48 hrs for short term toxicity study and up to 14 days for long term toxicity study after oral dose.

**Result:** Based on the mortalities and clinical signs observed in all female rats dosed with 2000 mg/kg body weight the LD50 value was calculated.

**Conclusion:** The study confirmed that the estimated LD50 value for methanolic extract of *Alpinia Officinarum* Hanse was 2000mg/kg body.

Keywords: Acute toxicity study, Alpinia Officinarum Hanse, OECD-425

Access this Article Online	Quick Response Code:
Website:http://www.journalofhospitalpharmacy.in	
Received on 30/10/2017	
Accepted on 07/11/2017 © PEN All rights reserved	国際786年2