

# ANTENATALLY DETECTED CONGENITAL CYSTIC ADENOMATOID MALFORMATION IN THE FETUS: SHOULD

## THE PREGNANCY BE CONTINUED? CASE SERIES AND REVIEW OF LITERATURE

*DrSnigdha Rao<sup>1</sup>, DrSnigdha Kumari<sup>2</sup>, DrAashima Arora<sup>3</sup>, Dr P.K. Saha<sup>4</sup>, DrRimpi Singla<sup>5</sup>*

DrSnigdhaKumari, Post Graduate Institute of Medical Education And Research, Chandigarh

*Address for Correspondence: editorjohp@gmail.com*

### ABSTRACT

Cystic adenomatoid malformations (CCAM) are developmental abnormalities of the lung which are considerably rare. They are benign hamartomas or dysplastic lung neoplasms with a decrease in the alveolar number and proliferation of terminal bronchioles. Prenatally detected CCAM has historically a poor outcome with perinatal mortality from ranging from 9% to 49%. But recent studies have shown a better prognosis of antenatally detected CCAM even if complicated by hydrops fetalis at presentation. This study presents two cases of fetuses with congenital cystic adenomatoid lung malformation and its management in pregnancy

#### *Access this Article Online*

Website:<http://www.journalofhospitalpharmacy.in> Quick Response Code:

Received on 29/05/2018  
Accepted on 30/05/2018 © HEB All rights reserved

