HEB



JOHP

Journal of Hospital Pharmacy

An Official Publication of Bureau for Health & Education Status Upliftment

(Constitutionally Entitled as Health-Education, Bureau)

Study of recovery of thrombocytopenia patients

Archana Puligari

Department of Pharmacy Practice & Pharm D.

Email Id: serviceheb@gmail.com

ABSTRACT

Background:

Thrombocytopenia occurs when the platelet count falls below the normal range value that is 1,50,000 cells / microliter of blood.

Objective:

To study the recovery of thrombocytopenia in different people.

Method:

A prospective study of thrombocytopenia cause, symptoms, diagnosis and treatment of a individual patient in a hospital.

Results:

Out of 20 patients, 10 patients recovered without a transfusion of the platelets with supportive treatment and ten patients recovered with transfusion of platelets when the platelet count falls below 20,000 or 30, 000cells\micro liter of blood volume and transfusion also depends on the severity of the patient's condition.

Conclusion:

Ten patients recovered with transfusion of Platelets when platelets count falls to 20,000 or 30, 000 cells\micro liter. Ten patients recovered without transfusion of platelets.

Keywords:

Thrombocytopenia, Platelets, Transfusion and Recovery.

Access this Article Online	Quick Response Code
Website: http://www.journalofhospitalpharmacy.in	
Received on 14/04/2023	
Accepted on 22/04/2023 © HEB All rights reserved	