

## A Review on Implication of Zika Virus Infection In Pregnancy

Dr. Purnima Ashok<sup>1</sup> and Dr. Sherin Sara Benny<sup>2</sup>

<sup>1</sup> Head of Department, KLE College of Pharmacy, Bengaluru.

<sup>2</sup> Assistant professor, KLE College of Pharmacy, Bengaluru.

**Address for Correspondence:** editorjohp@gmail.com

### ABSTRACT

Zika virus infection usually appears to be mild and self-limiting in most cases. But it may cause fetal abnormalities during pregnancy called Congenital Zika Syndrome, including Microcephaly. This teratogenic virus is transmitted through aedes species mosquito, mainly aedes albopictus. Pregnant women are more vulnerable to Zika virus at any time during gestation, especially first and second trimester which can be detected by using serum and urine real-time reverse transcription-polymerase chain reaction. Pregnant women should be advised to minimize mosquito bite and reduce sexual transmission risk by practicing safe sex.

**Key words:** Zika virus, aedes albopictus, pregnancy, Microcephaly, reverse transcription-polymerase chain reaction

#### *Access this Article Online*

Website: <http://www.journalofhospitalpharmacy.in> Quick Response Code:

Received on 01/02/2020  
Accepted on 10/03/2020 © HEB All rights reserved

